

Real-world characteristics, treatment patterns, and outcomes in patients with HR+/HER2– early breast cancer from China and the US

Pedram Razavi,¹ Qiao Li,² Mingxia Jiang,² Jiaxuan Liu,² Mengqi Zhang,² Hope S. Rugo,³ Joyce A. O'Shaughnessy,⁴ Erika Hamilton,⁵ Julia Kim,⁶ Murat Akdere,⁷ Baoling Sun,⁸ Vaidyanathan Ganapathy,⁹ Zuzanna Angehrn,⁷ Fen Ye,⁷ Binghe Xu²

¹Memorial Sloan Kettering Cancer Center and Weill Cornell Medical College, New York, NY, USA; ²National Cancer Center/National Clinical Research Center for Cancer/Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China; ³City of Hope Medical Centre, Duarte, CA, USA; ⁴Texas Oncology/Baylor University Medical Center, Dallas, TX, USA; ⁵Sarah Cannon Research Institute, Nashville, TN, USA; ⁶Genesis Research Group, Hoboken, NJ, USA; ⁷Novartis Pharma AG, Global Medical Affairs, Basel, Switzerland; ⁸Novartis China, Shanghai, China; ⁹Novartis Pharmaceuticals Corporation, East Hanover, NJ, USA

Disclaimer

- This material was reviewed by the Global Medical Review team and is recommended for external presentation in response to an unsolicited medical request subject to local approval
- This scientific information may include data/information on investigational uses of compounds/drugs that have not yet been approved by regulatory authorities
- Each CPO is responsible for ensuring that this material is used in accordance with local laws, regulations, industry, and Novartis codes and standards

Introduction (1 of 2)

- Previous studies have suggested that there are major differences between the US and China in the characteristics (eg, age, stage at diagnosis) and treatment patterns (eg, rates of [neo]adjuvant therapy, breast-conserving surgery) in patients with all subtypes of breast cancer (BC)^{1,2}
 - In the setting of hormone receptor positive/human epidermal growth factor receptor 2 negative (HR+/HER2-) early BC (EBC), the treatment landscape in China and the US is similar, with standard-of-care therapies including surgery, chemotherapy, and/or radiation therapy followed by adjuvant endocrine therapy^{3,4}
 - Further studies are needed to better understand the characteristics, treatment patterns, and outcomes in patients with HR+/HER2- EBC

Introduction (2 of 2)

- In the phase 3 NATALEE and monarchE trials, ribociclib and abemaciclib, respectively, demonstrated a reduction in the risk of recurrence in patients with HR+/HER2- EBC, which led to their approval in the adjuvant setting in both the US and China⁵⁻⁷
 - With the introduction of these treatments, assessment of treatment patterns and outcomes in Chinese and US patient populations can help in understanding unmet needs and aid in clinical decision-making in patients with high risk of recurrence
- In this analysis, we assess patient characteristics, eligibility for adjuvant CDK4/6 inhibitor, treatment patterns, and outcomes from Chinese and US electronic health record (EHR) databases of patients diagnosed with HR+/HER2- EBC between 2013 and 2022

Methods (1/2)

- This retrospective study used deidentified secondary data from the National Cancer Information Database (NCID) of China and the Flatiron Health US EHR database⁸
- Patients in this analysis were aged ≥ 18 and ≤ 80 years, were initially diagnosed with stage I to III HR+/HER2- EBC (in accordance with the American Joint Committee on *Cancer's Cancer Staging Manual*, 8th Edition) between 1 January 2013 and 31 December 2022, had undergone BC resection, and received ≥ 1 adjuvant therapy of any type (**Figure 1**)
 - Patients were identified as NATALEE or monarchE eligible based on criteria from each study (NATALEE: patients with N0 with high-risk features or node-positive [N1-3]; monarchE: patients N1 with high-risk features or N2-3)⁵⁻⁷
- Demographics and clinical characteristics were analyzed using descriptive statistics

Methods (2/2)

- Recurrence-free survival (RFS) and distant recurrence-free survival (DRFS) analyses, defined based on STEEP version 2.0 criteria, were conducted using the Kaplan-Meier method
 - RFS and DRFS analysis were assessed in the overall population and subgroups of patients defined by nodal status (N0 with high-risk features [N0-NATALEE-eligible], N0 without high-risk features [N0-NATALEE-ineligible], N1, and N2-3)

Results (1 of 6)

- In the China and US populations, 28,623 and 5930 patients met the selection criteria and were included in this analysis (**Figure 1**)
- The median duration of follow-up from initial diagnosis was 33.6 months (IQR, 19.2-56.4 months) in China and 52.3 months (IQR, 27.7-77.8 months) in the US

Figure 1. Flowchart of patient selection

Criteria	NCID China	Flatiron Health US
Initial EBC diagnosis from 1 Jan 2013 to 31 Dec 2022	N = 442,257 ^a	N = 12,138
Patients without other cancers at initial diagnosis	n = 401,871	n = 11,742 ^b
Patients diagnosed with HR+/HER2- EBC	n = 130,205	n = 6953
Patients aged ≥18 and ≤80 years at initial diagnosis	n = 129,222	n = 6451
Patients diagnosed with stage I-III disease	n = 104,394	n = 6197
Received primary tumor BC surgical resection at any time	n = 45,858	n = 6148
Received adjuvant therapy and had ≥2 visits after surgery	n = 28,623	n = 5930

^aIncludes patients diagnosed with EBC and metastatic BC. ^bIncludes patients without another malignancy any time prior to the EBC diagnosis. BC, breast cancer; EBC, early breast cancer; HER2-, human epidermal growth factor receptor 2-negative; HR+, hormone receptor positive; IQR, interquartile range; NCID, national cancer institute database.

Results (2 of 6)

Baseline characteristics and treatment patterns

- Characteristics and treatment patterns in patients in China and the US are included in **Table 1**
 - In the China and US populations, 16,743 (58.5%) and 2058 (34.7%) patients were NATALEE-eligible and 11,088 (38.7%) and 954 (16.1%) were monarchE-eligible, respectively
 - Patients in China vs the US were younger (median age 49 vs 63 years), were more likely to be premenopausal (44.6% vs 20.9%), and had higher rates of axillary node involvement (N0, 52.0% vs 71.5%; N1, 30.4% vs 22.1%; N2-3, 17.6% vs 6.4%), respectively
 - Patients in China vs the US were more likely to undergo total mastectomy (79.4% vs 37.1%) and receive adjuvant chemotherapy (79.0% vs 23.0%), respectively

Results (3 of 6)

Baseline characteristics and treatment patterns

Table 1. Demographics and Characteristics of Patients with HR+/HER2- EBC

Characteristic	NCID China (n = 28,623)	Flatiron Health US (n = 5930)	Anatomic stage, n (%)		
Median age (IQR), years	49 (43-58)	63 (54-70)	I	8666 (30.3)	3374 (56.9)
Sex, n (%)			II	14,149 (49.4)	2002 (33.8)
Female	28,467 (99.5)	5894 (99.4)	III	5808 (20.3)	554 (9.3)
Male	156 (0.5)	36 (0.6)	Type of surgery, n (%) ^b		
Menopausal status, n (%)			Mastectomy	22,725 (79.4)	2200 (37.1)
Pre-menopausal	12,761 (44.6)	1242 (20.9)	Breast conservation	2916 (10.2)	3961 (66.8)
Postmenopausal	9621 (33.6)	4374 (73.8)	Unknown	2982 (10.4)	-
Other (male patient)/unknown	6241 (21.8)	314 (5.3)	Type of adjuvant therapy, n (%) ^b		
Trial eligibility criteria, n (%)			Radiotherapy	18,337 (64.1)	3576 (60.3)
NATALEE-eligible	16,743 (58.5)	2058 (34.7)	Chemotherapy	22,623 (79.0)	1365 (23.0)
MonarchE-eligible	11,088 (38.7)	954 (16.1)	Endocrine therapy	25,071 (87.6)	5765 (97.2)
Nodal status, n (%)			CDK4/6 inhibitor	611 (2.1)	74 (1.2)
N0	14,886 (52.0)	4241 (71.5)	Other targeted therapy	-	3 (0.1) ^c
N0 NATALEE-ineligible	9474 (33.1)	3742 (63.1)	PARP inhibitor	-	4 (0.1)
N0 NATALEE-eligible	3010 (10.5)	499 (8.4)			
N0 risk unknown ^a	2402 (8.4)	0			
N1	8694 (30.4)	1311 (22.1)			
N2-3	5043 (17.6)	378 (6.4)			

^aThese patients had incomplete information to assess risk. ^bPatients may have received multiple treatment options; column percentages may add up to >100%. ^cFor the US cohort, other targeted therapy included alpelisib, everolimus, and idelalisib. CDK4/6, cyclin-dependent kinase 4 and 6; EBC, early breast cancer; HER2-, human epidermal growth factor receptor 2-negative; HR+, hormone receptor positive; IQR, interquartile range; N, nodal; PARP, poly(ADP-ribose) polymerase inhibitor.

Results (4 of 6)

Outcomes in China and US populations: RFS and DRFS

- 3- and 5-year RFS and DRFS rates were similar in the overall and overall NATALEE-eligible population in China and the US (Table 2)

Table 2. Outcomes in China and US: RFS and DRFS

	NCID China (n = 28,623)		Flatiron Health US (n = 5930)	
RFS (95% CI), %	3-year	5-year	3-year	5-year
Overall population	92.1 (91.7-92.5)	85.8 (85.2-86.4)	94.4 (93.7-95.0)	89.7 (88.7-90.6)
Overall NATALEE-eligible	89.1 (88.6-89.7)	81.2 (80.3-82.0)	89.2 (87.6-90.5)	81.8 (79.7-83.7)
DRFS (95% CI), %	3-year	5-year	3-year	5-year
Overall population	94.3 (94.0-94.7)	89.1 (88.5-89.6)	95.3 (94.6-95.8)	91.2 (90.3-92.0)
Overall NATALEE-eligible	91.7 (91.2-92.2)	84.5 (83.7-85.3)	90.4 (88.9-91.7)	83.6 (81.6-85.4)


CI, confidence interval; DRFS, distant recurrence-free survival; RFS, recurrence-free survival.

Results (5 of 6)

Outcomes in China and US Populations: RFS and DRFS

- Similar patterns were observed for RFS and DRFS by nodal status in China and US populations (Figures 2 and 3)

Figure 2. RFS by nodal status




DRFS, distant recurrence-free survival; NCID, national cancer institute database; RFS, recurrence-free survival.

Results (6 of 6)

Outcomes in China and US Populations: RFS and DRFS

Figure 3. DRFS by nodal status



DRFS, distant recurrence-free survival; NCID, national cancer institute database; RFS, recurrence-free survival.

Key Findings and Conclusions

- This harmonized analysis of 2 large, real-world databases highlights the differences in clinical characteristics and treatment patterns between patients with HR+/HER2- EBC treated in routine clinical practice in China and the US
- Patients with HR+/HER2- EBC in China were diagnosed at a younger age, were more likely to be premenopausal, and had higher rates of axillary node involvement, consequently they were more likely to undergo total mastectomy and be treated with adjuvant chemotherapy than patients in the US
- The risk of overall and distant recurrence was considerable in these 2 diverse populations, despite receiving available standard-of-care therapies
 - 3- and 5-year RFS and DRFS rates were similar both in China and the US in the overall population, across nodal status, and in the overall NATALEE-eligible population
- Despite differences in patient characteristics in China and the US, outcomes were comparable across these populations, and highlights a significant unmet need for patients with HR+/HER2- EBC

DRFS, distant recurrence-free survival; EBC, early breast cancer; HER2-, human epidermal growth factor receptor 2-negative; HR+, hormone receptor positive; RFS, recurrence-free survival.

References

1. Qian X, et al. *Transl Cancer Res.* 2023;12(7):1826-35.
2. Li J, et al. *JAMA Netw Open.* 2023;6(6):e2321388-e.
3. Huppert LA, et al. *CA Cancer J Clin.* 2023;73(5):480-515.
4. Harbeck N, Gnant M. *Lancet.* 2017;389(10074):1134-50.
5. Rastogi P, et al. *J Clin Oncol.* 2024;42(9):987-93.
6. Slamon DJ, *Ther Adv Med Oncol.* 2023;15:17588359231178125.
7. Fasching PA, et al. Poster presented at the ESMO Congress 2024; 13-17 September 2024. Barcelona, Spain.
8. Flatiron Health. Accessed 22 September 2025. <https://flatiron.com/databasecharacterization>.

Acknowledgements

The authors thank the patients enrolled in these studies and their families as well as the study investigators. Medical editorial assistance was provided by Nucleus Global and was funded by Novartis Pharmaceuticals Corporation. The authors had final responsibility for the poster

Disclosures

- **PR:** Institutional grant/funding support from AstraZeneca, Biodesix, Biothernostics, Biovica, Foresight Diagnostics, Grail, Guardant Health, Myriad, Novartis, Personalis, SAGA Diagnostics, Sophia Genetics, Tempus; and consultant/ad board/advisor fees from AstraZeneca, Foresight Diagnostics Guardant Health, Lilly, Myriad, Natera, NeoGenomics, Novartis, Pfizer, Prelude Therapeutics, Regor Pharmaceuticals, SAGA Diagnostics, and Tempus.
- **QL, MJ, JL, and MZ:** Nothing to disclose.
- **HR:** Grants from OBI Pharma, AstraZeneca, Pfizer, Novartis, Eli Lilly, Hoffmann-La Roche AG/Genentech, Merck, Daiichi Sankyo; personal fees from Daiichi Sankyo, Mylan/Viatris, NAPO, Eisai; grants from Gilead Sciences, Stemline Therapeutics, Ambrx.
- **JO:** Personal fees from Abbvie, Agendia, Amgen, Aptitude, AstraZeneca, Bristol-Myers Squibb, Celgene, Eisai, G1 Therapeutics, Genentech, Immunome, Ipsen Biopharmaceuticals, Lilly, Merck, Myriad, Novartis, Odonate Therapeutics, Pfizer, Puma, Prime, Roche, Seagen, Syndax, Carrick Therapeutics, Daiichi Sankyo, Gilead Sciences, Ontada, Pierre Fabre, Samsung, Sanofi.
- **EH:** Grants from Novartis, Hutchinson MediPharma, OncoMed, MedImmune, Stem CentRx, Curis, Verastem, Zymeworks, Syndax, Lycera, Rgenix, Millennium, TapImmune, Lilly, BerGenBio, Medivation, Kadmon, H3 Biomedicine, Radius Health, Acerta Pharma, Takeda, MacroGenics, Abbvie, Immunomedics, eFFECTOR Therapeutics, Mallinckrodt, Fujifilm, Merus, Nucana, TetraLogic Pharmaceuticals, PharmaMar, Regeneron, Leap Therapeutics, Taiho Pharmaceutical, Syros, CytomX, InventisBio, Oncothyreon; grants and other from Pfizer, Puma Biotechnology, Mersana, Boehringer Ingelheim, Cascadian Therapeutics; grants, nonfinancial support and other from Genentech/Roche, Lilly, Daiichi Sankyo, Eisai; grants and non-financial support from AstraZeneca, Tesaro, EMD Serono, Clovis, Novartis; personal fees from Flatiron; and non-financial support from Amgen, Bayer, Bristol-Myers Squibb, Genzyme, Helsinn Therapeutics, HERON, Lexicon, Medivation, Merck, Roche, Sysmex, Guardant Health, Foundation Medicine.
- **JK:** Personal fees from Genesis Research during the conduct of the study.
- **MA, BS, VG, ZA and FY:** Employment and stock ownership from Novartis. BX consulting or advisory role fees from AstraZeneca and Novartis.

Scan to obtain: Poster



<https://tinyurl.com/PedramRazavi438eP>

Copies of this poster obtained through Quick Response (QR) code are for personal use only and may not be reproduced without permission of the authors.

Presenting author email address: razavip@mskcc.org